

No Difference Between Red Wine or White Wine Consumption and Breast Cancer Risk

Polly A. Newcomb, PhD^{a,b} Hazel B. Nichols, MS^b, Jeannette M. Beasley, PhD, MPH^{a,c} Kathleen Egan, ScD^d, Linda Titus-Ernstoff, PhD^e, John M. Hampton, MS^b, Amy Trentham-Dietz, PhD^{b,f}

^aFred Hutchinson Cancer Research Center, Seattle, WA

^bUniversity of Wisconsin Paul P. Carbone Comprehensive Cancer Center, Madison, WI

^cGroup Health Cooperative, Seattle, WA

^dH. Lee Moffitt Cancer Center & Research Institute, Tampa, FL

^eDartmouth Medical School, Lebanon, NH

^fDepartment of Population Health Sciences, University of Wisconsin, Madison, WI

Corresponding Author:

Dr. Polly A. Newcomb

Fred Hutchinson Cancer Center, M4-B823

1100 Fairview Ave. N.

Seattle WA 98109-1024

Phone: 206-667-3476

Fax: 206-667-7850

Email: pnewcomb@fhcrc.org

Running title: Wine type and breast cancer risk

Key words: Breast cancer, alcohol, wine, case-control

Acknowledgements:

This study was supported by National Cancer Institute grants R01 CA47147, R01 CA47305, and R01 CA69664.

The authors are grateful to Drs. Henry Anderson, Patrick L. Remington, Meir J. Stampfer, Walter C. Willett, John A. Baron, E. Robert Greenberg and Gaia Pocobelli for advice at various stages of this study; Laura Stephenson and the staff of the Wisconsin Cancer Reporting System, Susan T. Gershman and the staff of the Massachusetts Tumor Registry; Marguerite Stevens and the staff of the New Hampshire Cancer Registry; and Linda Haskins, Heidi Judge, Laura Mignone, and Shafika Abrahams-Gessel along with the study interviewers and programmers in all three states for assistance with data collection. We are especially grateful to the study participants, whose generosity made this research possible.

Introduction

Epidemiologic studies have reported an increased risk of breast cancer among women who drink alcohol, including wine (1, 2) Two meta-analyses estimated a ~10% (95% CI 5%, 15%) increased risk of breast cancer with each additional 10g (~1 drink) of alcohol/day regardless of beverage type (3, 4). Few studies have evaluated breast cancer risk separately for red and white wine (5-8). There is some evidence of beneficial health effects of red wine from laboratory (9) and epidemiological studies of heart disease (10) and prostate cancer (11, 12). We evaluated overall alcohol as well as red and white wine consumption to examine beverage-specific effects on breast cancer.

Methods

The analysis was conducted in a population-based case-control study described fully elsewhere (13). Briefly, the study enrolled breast cancer cases identified from population-based registries in Wisconsin, Massachusetts (excluding metropolitan Boston), and New Hampshire, age 20-69 years, with incident invasive breast cancer diagnosed between 1995 and 2000. Controls were frequency matched to cases by 5-year age groups, selected from lists of licensed drivers (<65 years) or a roster of Medicare beneficiaries (\geq 65 years). Approximately 80% of cases (n=6,327) and 76% of controls (n=7,558) participated in the study; participation was similar in the three states.

Telephone interviews were administered concurrently for case (typically within one year of diagnosis) and control participants. Participants were queried on suspected risk factors for breast cancer, including age at menarche and menopause, age at first pregnancy, parity, postmenopausal hormone use, family history of breast cancer, smoking, education, and body mass index evaluated prior to the reference date. Alcohol use was evaluated as frequency of beer, red wine, white wine, or liquor (distilled spirits) consumption (number of drinks per day, week, or month) during the year previous to the reference date, defined as the date of diagnosis for cases or approximately the same period for controls. Logistic regression was used to

estimate odds ratios (OR) and 95% confidence intervals (CI). Variables were included in the multivariable model if statistically significantly associated with breast cancer ($P \leq 0.05$) in age-adjusted models. Models estimating the effect of wine type on risk were adjusted for all other sources of alcohol. To obtain P-values for trend, we included alcohol both as a continuous and categorical exposure. We tested for interaction by type of alcohol. Analyses were conducted using SAS (Version 9.1, SAS Institute, Cary, NC).

Results

Cases tended to have a younger age at menarche, later age at first birth, lower parity, and later age at menopause (data not shown). Current use of postmenopausal hormones, first-degree family history of breast cancer, greater body mass index, and higher education were also associated with increased breast cancer.

Any alcohol consumption in the year before the reference date was reported by similar proportions of cases (82%) and controls (82%). After multivariable adjustment, consumption of 14 or more drinks per week was associated with a 24% increase in breast cancer (95% CI: 3%-49%, P for trend= 0.003) compared to non-drinkers (Table 1). Menopausal status modified this association, with elevated risk observed only among postmenopausal women (P for interaction= 0.05). This effect was driven by a statistical interaction only among liquor drinkers. Higher intake of all beverage types was suggestive of increased risk, only elevated consumption of liquor was significantly elevated per drink (OR=1.02, 95% CI= 1.00 to 1.03). Wine consumption was not associated with risk of breast cancer (OR=1.01, 95% CI=0.99 to 1.02) and no differential was observed between red and white wine ($P=0.6$ for interaction, Table 2).

Discussion

In this large, multi-center, population-based case control study, alcohol was associated with an elevated risk of breast cancer, primarily because of increased risk in postmenopausal liquor drinkers. Neither red nor white wine was related to breast cancer.

The Women's Health Study, a randomized, chemopreventive trial of 39,876 health professionals followed for ~10 years, reported similar associations of white (multivariable RR 1.1, 95% CI 0.9-1.2) and red (multivariable RR 1.0, 95% CI 0.8-1.2) wine per 10 g/day increment on risk of breast cancer (6). Further, two other studies conducted in France reported findings that did not support inverse effects of red wine versus white wine. Hirvonen *et al* reported no significantly increased risk of breast cancer associated with either red or white wine after a median 6.6 years follow-up in a placebo-controlled antioxidant primary-prevention trial (8) with 95 breast cancer cases. A small case-control study of premenopausal women (n=154 cases and 154 controls) reported an increase in breast cancer for women who reported consuming more than 4 liters of red wine per month, or ~1 drink/day (OR=4.0; 95% CI 1.6-9.8; P-trend=0.003) (5). Whereas over half of the study population consumed red wine, <10% reported consuming white wine (5). This differs from our study where equal proportions of women reported consuming red and white wine. Another case-control study conducted in France reported a lower risk of breast cancer among women consuming 10 -12 g/d of wine (OR=0.51; 95% CI=0.30-0.97) when compared with non-wine drinkers, but they did not evaluate red and white wine separately(14).

Separate effects of red versus white wine on risk may have biological plausibility. Resveratrol, a polyphenol in grapes, peanuts, and other plant-based foods, has displayed anti-carcinogenic properties in mouse mammary cultures (15, 16). Concentrations are greater in red compared to white wine, due to the variable amounts of time that red grape skins are present during processing (17). However, resveratrol is metabolized quickly: consistent intake of two glasses of red wine daily would be equivalent to a dose of ~27 µg/kg body weight for a 70 kg individual, leading to detectable concentrations of derivatives but not free resveratrol (9). Thus these low levels may not result in bioeffective concentrations.

Limitations should be considered in interpreting findings. Exposure measurement was limited to recent use and patterns may have changed over time. Though we were able to control

for many confounders, we did not have data on drinking patterns (i.e. moderate versus binge drinking). Self-reported alcohol intake measures are prone to error due to both retrospective reporting and recall bias. However, alcohol intake measured via food frequency is correlated with food records (Spearman $r=0.9$) (18) and the reliability of reporting is acceptable ($r=0.8$) (19).

Wine consumption has been increasing over the past decade in the United States (20), so future studies may more precisely evaluate the effects red and white wine on incidence of breast cancer.

References

1. Smith-Warner SA, Spiegelman D, Yaun SS, et al. Alcohol and breast cancer in women: a pooled analysis of cohort studies. *Jama* 1998;279:535-540.
2. WCFR/AIRC. Alcoholic Drinks. In: Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global Perspective Washington DC: AICR; 2007. p. 157-171.
3. Ellison RC, Zhang Y, McLennan CE, Rothman KJ. Exploring the relation of alcohol consumption to risk of breast cancer. *Am J Epidemiol* 2001;154:740-747.
4. Key J, Hodgson S, Omar RZ, et al. Meta-analysis of studies of alcohol and breast cancer with consideration of the methodological issues. *Cancer Causes Control* 2006;17:759-770.
5. Viel JF, Perarnau JM, Challier B, Faivre-Nappez I. Alcoholic calories, red wine consumption and breast cancer among premenopausal women. *Eur J Epidemiol* 1997;13:639-643.
6. Zhang SM, Lee IM, Manson JE, et al. Alcohol consumption and breast cancer risk in the Women's Health Study. *Am J Epidemiol* 2007;165:667-676.
7. Vachon CM, Kushi LH, Cerhan JR, Kuni CC, Sellers TA. Association of diet and mammographic breast density in the Minnesota breast cancer family cohort. *Cancer Epidemiol Biomarkers Prev* 2000;9:151-160.
8. Hirvonen T, Mennen LI, de Bree A, et al. Consumption of antioxidant-rich beverages and risk for breast cancer in French women. *Ann Epidemiol* 2006;16:503-508.
9. Baur JA, Sinclair DA. Therapeutic potential of resveratrol: the in vivo evidence. *Nat Rev Drug Discov* 2006;5:493-506.
10. Klatsky AL, Friedman GD, Armstrong MA, Kipp H. Wine, liquor, beer, and mortality. *Am J Epidemiol* 2003;158:585-595.
11. Schoonen WM, Salinas CA, Kiemeneij LA, Stanford JL. Alcohol consumption and risk of prostate cancer in middle-aged men. *Int J Cancer* 2005;113:133-140.
12. Velicer CM, Kristal A, White E. Alcohol use and the risk of prostate cancer: results from the VITAL cohort study. *Nutr Cancer* 2006;56:50-56.

13. Sprague BL, Trentham-Dietz A, Newcomb PA, et al. Lifetime recreational and occupational physical activity and risk of in situ and invasive breast cancer. *Cancer Epidemiol Biomarkers Prev* 2007;16:236-243.
14. Bessaoud F, Daures JP. Patterns of alcohol (especially wine) consumption and breast cancer risk: a case-control study among a population in Southern France. *Ann Epidemiol* 2008;18:467-475.
15. Whitsett T, Carpenter M, Lamartiniere CA. Resveratrol, but not EGCG, in the diet suppresses DMBA-induced mammary cancer in rats. *J Carcinog* 2006;5:15.
16. Jang M, Cai L, Udeani GO, et al. Cancer chemopreventive activity of resveratrol, a natural product derived from grapes. *Science* 1997;275:218-220.
17. Kopp P. Resveratrol, a phytoestrogen found in red wine. A possible explanation for the conundrum of the 'French paradox'? *Eur J Endocrinol* 1998;138:619-620.
18. Giovannucci E, Colditz G, Stampfer MJ, et al. The assessment of alcohol consumption by a simple self-administered questionnaire. *Am J Epidemiol* 1991;133:810-817.
19. Longnecker MP, Newcomb PA, Mittendorf R, et al. The reliability of self-reported alcohol consumption in the remote past. *Epidemiology* 1992;3:535-539.
20. Wine Institute DoC. San Francisco: c 2005-2008 [cited 2007 Sept 11]. Available from: http://www.wineinstitute.org/industry/statistics/2007/wine_sales.php.

Table 1: Association of recent alcoholic beverage consumption and breast cancer risk

	All Women			Premenopausal Women			Postmenopausal Women ^{2,3}		
	Cases (n = 6,327)	Controls (n = 7,558)	OR ¹ (95% CI)	Cases (n = 2,462)	Controls (n = 2,925)	OR ¹ (95% CI)	Cases (n = 3,449)	Controls (n = 4,162)	OR ¹ (95% CI)
Drinks per week	No. (%)	No. (%)		No. (%)	No. (%)		No. (%)	No. (%)	
All alcohol									
Non-drinkers	1122 (18)	1379 (18)	1	351 (14)	424 (15)	1	709 (21)	876 (21)	1
<1	2129 (34)	2712 (36)	0.94 (0.85-1.04)	894 (36)	1045 (36)	1.02 (0.86-1.22)	1106 (32)	1480 (36)	0.90 (0.79-1.02)
1-3.4	1450 (23)	1731 (23)	0.99 (0.88-1.10)	592 (24)	767 (26)	0.90 (0.75-1.08)	745 (22)	863 (21)	1.03 (0.89-1.19)
3.5-6.9	702 (11)	793 (11)	1.02 (0.90-1.17)	306 (12)	363 (12)	0.96 (0.78-1.19)	345 (10)	384 (9)	1.05 (0.88-1.26)
7-13.9	619 (10)	658 (9)	1.11 (0.97-1.28)	221 (9)	223 (8)	1.17 (0.92-1.48)	355 (10)	395 (10)	1.07 (0.89-1.28)
≥ 14	305 (5)	285 (4)	1.24 (1.03-1.49)	98 (4)	103 (4)	1.10 (0.80-1.52)	189 (6)	164 (4)	1.37 (1.08-1.73)
P-trend			0.003			0.55			0.004
1-drink increase			1.01 (1.00-1.02)			1.00 (0.99-1.01)			1.02 (1.01-1.03)
Wine									
Non-drinkers	1122 (18)	1379 (18)	1	351 (14)	424 (15)	1	709 (21)	876 (21)	1
<1	2455 (39)	2920 (39)	1.03 (0.94-1.12)	1021 (42)	1183 (40)	1.09 (0.95-1.26)	1283 (37)	1550 (37)	0.97 (0.86-1.09)
1-3.4	1048 (17)	1248 (17)	1.00 (0.89-1.11)	424 (17)	546 (19)	0.98 (0.82-1.17)	538 (16)	634 (15)	0.95 (0.81-1.11)
3.5-6.9	343 (5)	363 (5)	1.13 (0.95-1.34)	134 (5)	149 (5)	1.15 (0.87-1.50)	179 (5)	192 (5)	1.05 (0.83-1.33)
7-13.9	256 (4)	299 (4)	1.07 (0.89-1.29)	90 (4)	86 (3)	1.42 (1.02-1.96)	147 (4)	192 (5)	0.93 (0.73-1.19)
≥ 14	82 (1)	70 (1)	1.37 (0.98-1.91)	29 (1)	24 (1)	1.58 (0.90-2.79)	47 (1)	43 (1)	1.23 (0.80-1.90)
P-trend			0.20			0.12			0.84
1-drink increase			1.01 (0.99-1.02)			1.00 (0.99-1.02)			1.01 (0.99-1.03)
Beer									
Non-drinkers	1122 (18)	1379 (18)	1	351 (14)	424 (15)	1	709 (21)	876 (21)	1
<1	1756 (28)	2040 (27)	1.01 (0.93-1.10)	714 (29)	870 (30)	0.92 (0.80-1.05)	916 (27)	1044 (25)	1.05 (0.94-1.18)
1-3.4	754 (12)	836 (11)	1.06 (0.95-1.20)	359 (15)	444 (15)	0.91 (0.77-1.08)	345 (10)	346 (8)	1.21 (1.02-1.44)
3.5-6.9	163 (3)	210 (3)	0.90 (0.72-1.12)	105 (4)	120 (4)	0.91 (0.69-1.21)	48 (1)	76 (2)	0.81 (0.55-1.17)
7-13.9	129 (2)	139 (2)	1.08 (0.84-1.39)	52 (2)	55 (2)	1.05 (0.71-1.56)	66 (2)	74 (2)	1.08 (0.76-1.52)
≥ 14	82 (1)	84 (1)	1.12 (0.81-1.53)	32 (1)	39 (1)	0.89 (0.55-1.45)	45 (1)	39 (1)	1.44 (0.92-2.25)
P-trend			0.55			0.28			0.11
1-drink increase			1.01 (1.00-1.02)			1.01 (0.99-1.02)			1.01 (1.00-1.03)
Liquor									
Non-drinkers	1122 (18)	1379 (18)	1	351 (14)	424 (15)	1	709 (21)	876 (21)	1
<1	2463 (39)	3132 (41)	0.90 (0.83-0.98)	1098 (45)	1371 (47)	0.90 (0.79-1.02)	1186 (34)	1529 (37)	0.93 (0.83-1.04)
1-3.4	940 (15)	1044(14)	1.00 (0.90-1.12)	334 (14)	402 (14)	0.92 (0.77-1.11)	541(16)	582 (14)	1.07 (0.92-1.24)
3.5-6.9	154 (2)	146 (2)	1.14 (0.90-1.44)	55 (2)	61 (2)	0.93 (0.63-1.37)	93 (3)	75 (2)	1.40 (1.02-1.93)
7-13.9	162 (3)	175 (2)	1.00 (0.80-1.26)	37 (2)	38 (1)	1.00 (0.62-1.60)	115 (3)	131 (3)	0.98 (0.75-1.29)
≥ 14	95 (2)	73 (1)	1.41 (1.03-1.94)	18 (1)	19 (1)	1.10 (0.56-2.13)	74 (2)	50 (1)	1.63 (1.12-2.38)
P-trend			0.28			0.37			0.04
1-drink increase			1.02 (1.00-1.03)			0.99 (0.97-1.02)			1.03 (1.01-1.05)

¹ Adjusted for age, state, family history of breast cancer, age at menarche, age at first birth, parity, menopausal status, age at menopause, postmenopausal hormone use, body mass index, education, wine, beer and liquor consumption.

² For the test of the interaction between total alcohol consumption (continuous) and menopausal status P = 0.05.

³ For the test of the interaction between alcohol consumption (continuous) and menopausal status: all alcohol P=0.05; wine P=0.89, beer P=0.89, liquor P=0.03

Table 2: Association of recent red and white wine consumption and breast cancer risk

Wine type	Cases (n= 6,327)	Controls (n= 7,558)	OR (95% CI) ^{1,2}
Drinks per week	No. (%)	No. (%)	
Red wine			
Non-drinkers	1,122 (18)	1,379 (18)	1
<1	2,145 (34)	2,502 (33)	1.03 (0.94-1.12)
1-3.4	738 (12)	885 (12)	0.94 (0.83-1.06)
3.5-6.9	98 (2)	119 (2)	0.91 (0.69-1.21)
7-13.9	124 (2)	133 (2)	1.17 (0.90-1.51)
≥14	28 (<1)	24 (<1)	1.27 (0.73-2.22)
P-trend			0.99
1-drink increase			1.00 (0.98-1.01)
White wine			
Non-drinkers	1,122 (18)	1,379 (18)	1
<1	2,145 (34)	2,587 (34)	0.98 (0.90-1.07)
1-3.4	880 (14)	997 (13)	1.06 (0.94-1.19)
3.5-6.9	161 (3)	143 (2)	1.40 (1.10-1.78)
7-13.9	103 (2)	137 (2)	0.90 (0.69-1.18)
≥14	40 (1)	34 (<1)	1.37 (0.86-2.19)
P-trend			0.34
1-drink increase			1.02 (1.00-1.04)

¹ Adjusted for age, state, family history of breast cancer, age at menarche, age at first birth, parity, menopausal status, age at menopause, postmenopausal hormone use, body mass index, education, total alcohol consumption (minus red or white wine consumption).

² For the test of the interaction wine type (red vs. white, continuous) P=0.60.