

Supplementary Table 1. Descriptive factors in the non WT1-syndromic cohort

	ESRD Status		PBWT		CRF		All
	N	%	N	%	N	%	N
Bilateral WT Status							
Synchronous bilateral	469	6	29	64.4	14	25.5	512
Metachronous bilateral	70	0.9	16	35.6	1	1.8	87
Unilateral	7310	93.1	0	0	40	72.7	7350
Sex							
Male	3505	44.7	18	40	18	32.7	3541
Female	4344	55.3	27	60	37	67.3	4408
Ethnicity							
White	5593	71.3	31	68.9	37	67.3	5661
Black	1386	17.7	7	15.6	11	20.0	1404
Hispanic	682	8.7	6	13.3	4	7.3	692
Other	188	2.4	1	2.2	3	5.5	192
NWTS Study Cohort							
I	451	5.7	3	6.7	10	18.2	464
II	642	8.2	2	4.4	9	16.4	653
III	1858	23.7	12	26.7	17	30.9	1887
IV	2495	31.8	18	40	15	27.3	2528
V	2403	30.6	10	22.2	4	7.3	2417

PBWT: progressive bilateral Wilms tumor; CRF: chronic renal failure.

%

6.4

1.1

92.5

44.5

55.5

71.2

17.7

8.7

2.4

5.8

8.2

23.7

31.8

30.4